

JOURNAL OF ALGEBRA 112, 139–150 (1988)

Flatness, LCM-Stability, and Related Module-Theoretic Properties

D. D. ANDERSON*

*Department of Mathematics, University of Iowa,
Iowa City, Iowa 52242*

AND

DAVID E. DOBBS

*Department of Mathematics, University of Tennessee,
Knoxville, Tennessee 37996**Communicated by J. Dieudonné*

Received June 5, 1986

The relationship between flatness and LCM-stability is clarified by the following two results. A finitely generated ideal I of an integral domain is flat if and only if I is n -flat for some integer $n \geq 2$. There exists an integral domain with a nonflat ideal $J = (a, b, c)$ such that $Jab \cap Jac \cap Jbc = J(ab, ac, bc)$. Next, related module-theoretic properties, $(**)$ and $(*)$, respectively weaker than projectivity and flatness, are introduced. Under appropriate finiteness conditions, these properties are preserved by certain inverse limits. Their study leads to new characterizations of quasi-complete local rings and coherent integral domains. © 1988 Academic Press, Inc.

1. INTRODUCTION

This article is a sequel to [2, 6]. All rings are assumed commutative and all modules unital.

Sections 2 and 3 investigate further the relationship between flatness and LCM-stability. Recall that a torsion-free module E over an integral domain R is said to be LCM-stable over R in case $E(Ra \cap Rb) = Ea \cap Eb$ for all $a, b \in R$. (This property was first studied in [7, 12] in case E is an extension domain of R .) Although flatness implies LCM-stability, Uda [12] has shown that the converse is false. However, [6, Corollary 2.3] established that, for 2-generated ideals I of an integral domain R , a certain weak version of LCM-stability is actually equivalent to flatness (and hence, if $I \neq 0$, to invertibility). Our first result, Example 2.1, shows that there is no

* D.D.A. acknowledges the support services provided at University House, The University of Iowa.

analogous characterization of flatness for 3-generated ideals. On a positive note, Theorem 3.5 establishes that, for finitely generated ideals of an integral domain, LCM-stability is equivalent to flatness.

Indeed, more is shown. First, recall that a module E over an integral domain R is called n -flat (over R) in case each linear relation of n elements $r_1, \dots, r_n \in R$ with coefficients in E is a linear consequence of linear relations of the r_i 's with coefficients in R . It is known that flatness is equivalent to " n -flat for each positive integer $n \geq 2$ " and that 2-flatness is equivalent to LCM-stability [6, Theorem 3.3(b)]. Now, we can state a sharper form of Theorem 3.5: a finitely generated ideal I of an integral domain R is R -flat if (and only if) I is n -flat over R for some positive integer $n \geq 2$. Also of some independent interest in Section 3 is Proposition 3.1, which connects n -flatness and tensor product.

Sections 4 and 5 take their inspiration from the characterizations in [2] of invertible and flat ideals. This work had been the principal motivation for the study of LCM-stability in [6]. We abstract these characterizing properties to the module-theoretic setting, obtaining properties called $(**)$ and $(*)$. (See Section 4 for their definitions.) It turns out that $(**)$ (resp., $(*)$) is implied by, but is not equivalent to, projectivity (resp., flatness). The main result of Section 4, Theorem 4.5, states that if R is a local ring, then its completion \hat{R} satisfies $(**)$ as an R -module if and only if R is quasi-complete (in the sense of [8, 1]). The main result of Section 5 is that, under appropriate finiteness conditions but not in general, $(**)$ and $(*)$ are each preserved by arbitrary products. (See Theorem 5.3 and Proposition 5.1(c).) Section 5 also includes a new $(*)$ -theoretic characterization of coherent integral domains (in Corollary 5.6) and new proofs of some known properties of such integral domains.

2. A NONCRITERION FOR FLATNESS

It was shown in [6, Corollary 2.3] that a 2-generated ideal $I = Ra + Rb$ of an integral domain R is R -flat if (and only if) $Ia \cap Ib = I(Ra \cap Rb)$. We show next that there is no analogous criterion for flatness of 3-generated ideals.

EXAMPLE 2.1. Let X, Y, Z be commuting, algebraically independent indeterminates over a field k . Set $R = k[X, Y, Z]$ and $I = RXY + RXZ + RYZ$. Then

$$IXY \cap IXZ \cap IYZ = I(RXY \cap RXZ \cap RYZ)$$

although I is not R -flat.

Proof. We claim that $I^{-1} = R$. Indeed, if $\delta \in I^{-1}$, write $\delta = fg^{-1}$ for relatively prime polynomials f, g in the UFD R . As $fI \subset gR$, it follows that

g is a factor of each of XY , XZ , and YZ , whence g is a constant. Thus $\delta \in R$, as claimed. Hence $II^{-1} = I \neq R$, so that I is not invertible. Alternatively, notice that I , though finitely generated, is not principal. As R is factorial, it follows that I is not invertible. Being finitely generated and non-zero, I is therefore not flat.

Put $J = IXY \cap IXZ \cap IYZ$ and $K = I(RXY \cap RXZ \cap RYZ)$. Evidently $K \subset J$. We shall show $J \subset K$. View J as

$$J = (X^2Y^2, X^2YZ, XY^2Z) \cap (X^2YZ, X^2Z^2, XYZ^2) \cap (XY^2Z, XYZ^2, Y^2Z^2).$$

As $K = IXYZ$, it is possible, via a long but straightforward argument exploiting the factoriality of R , to show $J \subset K$. We leave the details for the reader. For a more elegant argument, one may use the fact that X, Y, Z is an R -sequence to conclude that J is generated by monomials of the form $X^\alpha Y^\beta Z^\gamma$, where $\alpha, \beta, \gamma \geq 1$ and at most one of α, β, γ is 1. As $X^2Y^2Z, X^2YZ^2, XY^2Z^2 \in J$ it follows that each such $X^\alpha Y^\beta Z^\gamma$ is in K , whence $J \subset K$. ■

3. FINITELY GENERATED LCM-STABLE IDEALS ARE FLAT

The following useful result connects n -flatness and tensor products.

PROPOSITION 3.1. *Let M be an n -flat module over an integral domain R , for some positive integer n . If I is an n -generated ideal of R , then the canonical homomorphism $g: I \otimes_R M \rightarrow M$ is a monomorphism.*

Proof. We shall show that each $e \in \ker(g)$ is 0. To this end, write $I = Ri_1 + \cdots + Ri_n$, and note that $e = \sum i_j \otimes m_j$, for suitable $m_1, \dots, m_n \in M$ such that $\sum i_j m_j = g(e) = 0$. As M is n -flat over R , there exist elements $r_{kj} \in R$ and $m_j^* \in M$ such that

$$m_k = \sum_j r_{kj} m_j^* \quad \text{for } k = 1, \dots, n$$

and

$$\sum_{k=1}^n i_k r_{kj} = 0 \quad \text{for each } j.$$

Then

$$\begin{aligned} e &= \sum_k i_k \otimes \left(\sum_j r_{kj} m_j^* \right) = \sum_j \left(\sum_k i_k r_{kj} \otimes m_j^* \right) \\ &= \sum 0 \otimes m_j^* = \sum 0 = 0. \quad \blacksquare \end{aligned}$$

Remark 3.2. Calculations similar to the one above are given, for $n = 2$, in the proofs of [6, Corollary 3.4 and Theorem 3.8]. Moreover, with the

aid of a standard criterion for flatness [3, Proposition 1, p. 37], one sees easily that Proposition 3.1 leads to a new proof of [6, Corollary 3.4].

We next infer a companion for [6, Theorem 2.2].

COROLLARY 3.3. *Let I be an n -generated ideal of an integral domain R , for some positive integer n . Then I is R -flat if (and only if) I is n -flat over R .*

Proof. As I is n -flat, Proposition 3.1 assures that the multiplication map $I \otimes_R I \rightarrow I$ is a monomorphism. Then, by [5, Proposition 1], I is R -flat. ■

Remark 3.4. The preceding result is best-possible. Indeed, a 1-flat 2-generated ideal I of an integral domain R need not be R -flat. To see this, observe that if an integral domain R is not a Prüfer domain, then some (necessarily 1-flat) 2-generated ideal of R is not R -flat.

The main result of this section is

THEOREM 3.5. *Let I be a finitely generated ideal of an integral domain R . Then the following conditions are equivalent:*

- (1) I is LCM-stable over R ;
- (2) I is n -flat over R for some positive integer $n \geq 2$;
- (3) the canonical homomorphism $I \otimes_R J \rightarrow J$ is a monomorphism for each 2-generated ideal J of R such that $J \subset I$;
- (4) I is R -flat.

Proof. (4) \Rightarrow (2): Trivial.

(2) \Rightarrow (1): It is straightforward to verify that if $m < n$ are positive integers and a module E is n -flat over R , then E is m -flat over R . As LCM-stability is equivalent to 2-flatness, the assertion follows.

(1) \Rightarrow (3): By (1), I is 2-flat over R and so, by Proposition 3.1, $J \otimes_R I \rightarrow JI$ is monomorphism. As $J \otimes_R I \cong I \otimes_R J$, the assertion follows.

(3) \Rightarrow (4): Without loss of generality, R is quasi-local. It is enough to show that I is a principal ideal of R . Deny. Consider a minimal generating set $S = \{a, b, \dots\}$ for I as an ideal of R , with $a \neq b$. Set $J = Ra + Rb$. By minimality, $v(J) = 2$. (As usual, $v(E)$ denotes the minimum cardinality of a generating set for an R -module E .) Moreover, by (3), $I \otimes_R J \cong IJ$, and so $v(I \otimes_R J) = v(IJ)$. As $v(I \otimes_R J) = v(I) v(J)$, it follows that

$$\begin{aligned} 2v(I) &= v(Ia + Ib) = |\{xy : x \in S, y \in \{a, b\}\}| \\ &\leq 2|S| - 1 \quad (\text{since } ab = ba) \\ &= 2v(I) - 1, \end{aligned}$$

the desired contradiction. ■

Remark 3.6. (a) For future applications, it may be helpful to make the following observation, in the spirit of [5, Proposition 2]. Formally as before, one can define what is meant by n -flatness of a module over an arbitrary (commutative) ring. Then, by the above proof that (3) \Rightarrow (4), we have the following result. If I is a finitely generated 2-flat ideal of a (commutative) ring R , then I is locally principal over R .

(b) The implication (3) \Rightarrow (4) in Theorem 3.5 establishes the converse of Proposition 3.1 in case M is a finitely generated ideal of R and $n = 2$. It would be interesting to have additional instances of the converse of Proposition 3.1.

4. ON INTERSECTIONS AND QUASI-COMPLETENESS

This section treats the module-theoretic properties, dubbed (*) and (**) in the following, which underlay the ideal-theoretic studies in [2]. Let R be a ring and let E be an R -module. We shall say that E satisfies (*) (over R) if $(I \cap J)E = IE \cap JE$ for all ideals I, J of R , and that E satisfies (**) (over R) if $(\bigcap I_\alpha)E = \bigcap (I_\alpha E)$ for all families $\{I_\alpha\}$ of ideals of R . By adapting the proofs of [2, Theorems 2 and 1], we easily have

PROPOSITION 4.1. *Let R be an integral domain and E a torsion-free R -module. Then:*

(a) *E satisfies (*) if and only if E is R -flat.*

(b) *Suppose, in addition, that E is a fractional ideal of R . Then E satisfies (**) if and only if E is R -projective.*

Remark 4.2. The above properties are related in general as follows. If R is a ring and E is an R -module, then

$$E \text{ is } R\text{-projective} \Rightarrow E \text{ satisfies } (**) \text{ over } R$$

$$\Downarrow \qquad \qquad \Downarrow$$

$$E \text{ is } R\text{-flat} \Rightarrow E \text{ satisfies } (*) \text{ over } R.$$

Proof. The “vertical” implications are clear. Moreover, as in the proof of [2, Theorem 2, (3) \Rightarrow (1)], it follows via [3, Proposition 6, p. 17] that flatness implies (*). Finally, to show that each projective R -module E satisfies (**), choose P so that $E \oplus P$ is a free R -module F . Then

$$\bigcap (I_\alpha E) \oplus \bigcap (I_\alpha P) = \bigcap (I_\alpha E \oplus I_\alpha P) = \bigcap (I_\alpha F).$$

Expressing F in terms of an R -basis shows that this intersection is just $(\bigcap I_\alpha)F$, and hence is just $(\bigcap I_\alpha)E \oplus (\bigcap I_\alpha)P$. Equating first summands as submodules of E gives $\bigcap (I_\alpha E) = (\bigcap I_\alpha)E$, completing the proof. ■

The above result has natural applications. For instance, if a commutative R -algebra S is free as an R -module, then S satisfies $(**)$ over R . In particular, for each positive integer n , the polynomial ring $R[X_1, \dots, X_n]$ satisfies $(**)$ over R . The same holds for the formal power series ring $R[[X]]$ if R is Noetherian: this will follow from Theorem 5.3(b) below.

By Proposition 4.1(a), an overring S of an integral domain R satisfies $(*)$ over R if and only if S is R -flat. We show next that the situation for $(**)$ is very different.

THEOREM 4.3. *If an overring S of an integral domain R satisfies $(**)$ over R , then $S = R$.*

Proof. Assume that S satisfies $(**)$. Then, by Remark 4.2 and Proposition 4.1(a), S is R -flat. Thus, by [11, Lemma 1 and Theorem 1], $(b:a)S = S$ for all nonzero $a, b \in R$ such that $ab^{-1} \in S$. (As usual, $(b:a)$ denotes $\{r \in R: ra \in Rb\}$.) Let $I = \bigcap (b:a)$, the intersection being indexed by the elements $ab^{-1} \in S$. By hypothesis, $IS = \bigcap ((b:a)S)$, which is just $\bigcap S = S$. In particular, $I \neq 0$; pick a nonzero element c of I . Now, for each $ab^{-1} \in S$, $c \in (b:a)$, whence $ab^{-1} \in Rc^{-1}$; thus $S \subset Rc^{-1}$. As S is then a fractional ideal of R satisfying $(**)$, Proposition 4.1(b) yields that S is invertible. Hence S is finitely generated as an R -module. It follows that S is integral over R . However, there are no flat integral proper overrings [11, Proposition 2], so $S = R$, completing the proof. ■

Despite the preceding result, some important algebras (which need not be free as modules) *do* satisfy $(**)$. We shall show this in Examples 4.4 and Theorem 4.5. To motivate that work, consider the above concepts for a special R . It is easy to see that if R is a valuation domain (more generally, a ring whose ideals are linearly ordered by inclusion), then *each* R -module E satisfies $(*)$; and that if R is a DVR with local uniformizing parameter π , then an R -module E satisfies $(**)$ if and only if $\bigcap \pi^n E = 0$. These considerations lead naturally to the DVR $\mathbb{Z}_{p\mathbb{Z}}$, which appears next in Example 4.4(a).

The following examples give extra motivation for Theorems 4.5 and 5.3(b).

EXAMPLES 4.4. (a) Despite Proposition 4.1(b) and Remark 4.2, $(**)$ does not imply projectivity, even for a torsion-free module over a DVR. For instance, let p be a prime positive integer, let $R = \mathbb{Z}_{p\mathbb{Z}}$, and let $E = \hat{R}$. View E as the integral domain \mathbb{Z}_p of p -adic integers. It can be shown that E satisfies $(**)$ over R . (To see this, one may apply the above comment to the DVR $\mathbb{Z}_{p\mathbb{Z}}$, since $\bigcap p^n \hat{\mathbb{Z}}_{p\mathbb{Z}} = 0$; or one may appeal to Theorem 4.5.) As R may be viewed as a subdomain of E , E is a torsion-free R -module; however, E is not free as an R -module (cf. [9, Theorem 18]).

(b) The following additional example that “(**) $\not\Rightarrow$ projectivity” will be generalized in Section 5. Let $R = \mathbb{Z}$ and let $E = \prod \mathbb{Z}$, the product of denumerably many copies of \mathbb{Z} . It can be shown that E satisfies (**) over R (via a direct calculation using the definition of (**) and standard facts about \mathbb{Z} , or via Theorem 5.3(b)). Of course, E is torsion-free over R ; however, by a celebrated result of Baer (cf. [9]), E is not a free R -module. ■

To set the stage for a generalization of Example 4.4(a), consider (R, M) , a local (Noetherian) ring R with maximal ideal M . Let (\hat{R}, \hat{M}) denote the M -adic completion of R . It is well known that \hat{R} is R -flat; hence, by Remark 4.2, \hat{R} satisfies (*) over R . Before determining when \hat{R} satisfies (**) over R , it is convenient to recall the following material (cf. [8, 1]). We say that (R, M) is *quasi-complete* if, for each descending sequence $B_1 \supset B_2 \supset \dots$ of ideals of R and each integer k , there exists an integer $m(k)$ such that $B_{m(k)} \subset (\cap B_n) + M^k$. Equivalently, R is quasi-complete if the assignment $I \mapsto I\hat{R}$ gives a lattice-isomorphism $L(R) \rightarrow L(\hat{R})$ from the lattice $L(R)$ of ideals of R to the lattice $L(\hat{R})$ of ideals of \hat{R} . Completeness of R (that is, the condition $R \cong \hat{R}$) implies quasi-completeness, but the converse is false. Indeed, if R is a one-dimensional (still local) integral domain, then R is quasi-complete if and only if \hat{R} is an integral domain. Thus, for instance, $\mathbb{Z}_{p\mathbb{Z}}$ and $\mathbb{Q}[X]_{(X)}$ are each quasi-complete but not complete.

The main result of this section is

THEOREM 4.5. *Let (R, M) be a local ring. Then \hat{R} satisfies (**) over R if and only if R is quasi-complete.*

Proof. The “if” part follows since the (complete) lattice-isomorphism $L(R) \rightarrow L(\hat{R})$ preserves arbitrary intersections. Conversely, let \hat{R} satisfy (**). Consider a descending sequence $\{B_n\}$ of ideals of R and let k be a positive integer. Since \hat{R} is (quasi-) complete, there exists a positive integer $m(k)$ such that

$$B_{m(k)}\hat{R} \subset (\cap (B_n\hat{R}) + \hat{M}^k.$$

However, $\cap (B_n\hat{R}) = (\cap B_n)\hat{R}$ since \hat{R} satisfies (**). Using standard facts about completions (cf. [10, Proposition 6, p. 95]), we find

$$\begin{aligned} B_{m(k)} &= B_{m(k)}\hat{R} \cap R \subset \left(\left(\cap B_n \right) \hat{R} + \hat{M}^k \right) \cap R \\ &= \left(\left(\cap B_n \right) \hat{R} + M^k \hat{R} \right) \cap R = \left(\left(\left(\cap B_n \right) + M^k \right) \hat{R} \right) \cap R \\ &= \left(\cap B_n \right) + M^k. \end{aligned}$$

Hence R is quasi-complete, as asserted. ■

5. ON DIRECT PRODUCTS AND COHERENCE

Motivated in part by Example 4.4(b), we devote this section to studying when a product of modules satisfies (**). By also considering the analogous question for (*), we shall find that applications to coherent rings arise naturally.

It is convenient next to recall the following material from [4, Theorems 2.1 and 2.2]. A ring R is said to be *coherent* in case each finitely generated ideal of R is finitely presented. (The most familiar examples of coherent rings are arbitrary Noetherian rings and arbitrary Prüfer domains.) Equivalently, R is coherent in case each of the following three equivalent conditions holds: if $\{M_\beta\}$ is a set of flat R -modules, then $\prod M_\beta$ is R -flat; each product of (arbitrarily many) copies of R is R -flat; $I \cap J$ is finitely generated for each pair of finitely generated ideals I and J of R , and the annihilator $(0:a) = \{r \in R: ra = 0\}$ is finitely generated for each $a \in R$. This section's techniques will lead to new proofs of some of these facts in case R is an integral domain, as well as giving a new characterization of coherent integral domains (see Corollary 5.6).

The next result collects some basic information.

PROPOSITION 5.1. *Let R be a ring and $\{E_\beta\}$ a set of R -modules. Set $S = \bigoplus E_\beta$ and $P = \prod E_\beta$. Then:*

- (a) *S satisfies (*) (resp., (**)) if and only if E_β satisfies (*) (resp., (**)) for each β .*
- (b) *If P satisfies (*) (resp., (**)), then E_β satisfies (*) (resp., (**)) for each β .*
- (c) *If E_β satisfies (**) (or (*)) for each β , then P need not satisfy (*) (or (**)).*

Proof. (a) For each ideal I of R , $IS = I \sum E_\beta = \sum IE_\beta = \bigoplus IE_\beta$. Thus, for ideals I_α of R , one has

$$\left(\bigcap I_\alpha\right)S = \bigoplus_\beta \left(\left(\bigcap_\alpha I_\alpha\right)E_\beta\right)$$

and

$$\bigcap (I_\alpha S) = \bigcap \left(\bigoplus_\beta I_\alpha E_\beta\right) = \bigoplus_\beta \left(\bigcap_\alpha (I_\alpha E_\beta)\right).$$

Thus, $(\bigcap I_\alpha)S = \bigcap (I_\alpha S)$ if and only if $(\bigcap I_\alpha)E_\beta = \bigcap (I_\alpha E_\beta)$ for each β . The assertions follow easily.

(b) Apply (a), noting that $P = E_\beta \oplus (\prod \{E_\gamma; \gamma \neq \beta\})$.

(c) Consider any integral domain R which is not coherent. By the above-cited work of Chase, some product $P = \prod R$ of copies of R is not R -flat. As P is torsion-free over R , Proposition 4.1(a) and Remark 4.2 yield that P does not satisfy (*) and, hence, that P does not satisfy (**). However (each copy of) R is R -projective, and hence satisfies both (**) and (*). The proof is complete. ■

We shall also require the following result, which is rather well known.

LEMMA 5.2. *Let R be a ring and I an ideal of R such that $v(I) = \gamma$. Then the following conditions are equivalent:*

(1) *I is finitely generated.*

(2) *For each collection $\{E_\beta\}$ of R -modules, the canonical inclusion map $I \prod E_\beta \rightarrow \prod IE_\beta$ (of submodules of $\prod E_\beta$) is the identity.*

(3) *There exists an indexed family $\{E_\beta\}$ of copies of R (that is, $E_\beta = R$ for each β), with index set having cardinality at least γ , such that $I \prod E_\beta = \prod (IE_\beta)$ (that is, such that $I \prod R = \prod I$).*

Theorem 5.3 is the main result in this section. It is important for at least three reasons. First, it reduces the assertion (in Example 4.4(b)) that $\mathbb{Z} \times \mathbb{Z} \times \cdots$ satisfies (**) over \mathbb{Z} to the obvious assertion that \mathbb{Z} (is Noetherian and) satisfies (**) over itself. Second, Theorem 5.3(b) implies, as promised prior to Theorem 4.3, that if R is a Noetherian ring, then $R[[X]]$ satisfies (**) over R : one need only note that $R[[X]] \cong R \times R \times \cdots$ as R -modules and that R satisfies (**) over itself. Third, the result is sharp, for both its parts would fail without their “coherent” and “Noetherian” hypotheses: see the example in Proposition 5.1(c).

THEOREM 5.3. *Let R be a ring, $\{E_\beta\}$ a collection of R -modules, and $P = \prod E_\beta$. Then:*

(a) *If R is coherent and if E_β satisfies (*) over R for each β , then P satisfies (*) over R .*

(b) *If R is Noetherian and if E_β satisfies (**) over R for each β , then P satisfies (**) over R .*

Proof. We begin with a useful observation: in the definition of “satisfies (*)” (resp., (**)), the test ideals I, J (resp., I_α) may be assumed finitely generated. For such data, $I \cap J$ is finitely generated in part (a) by coherence, and $\cap I_\alpha$ is finitely generated in part (b) since R is Noetherian. A common proof for (a) and (b) can now be given, in terms of an indexed family $\{I_\gamma\}$ of finitely generated ideals, viewed as $\{I, J\}$ in part (a) and as $\{I_\alpha\}$ in part (b). By combining the above remarks about finite generation,

condition (2) in Lemma 5.2, the hypotheses on E_β , and commutativity of \cap and \prod , we have

$$\left(\bigcap I_\gamma\right)P = \prod_\beta \left(\left(\bigcap_\gamma I_\gamma\right)E_\beta\right) = \prod_\beta \left(\bigcap_\gamma (I_\gamma E_\beta)\right) = \bigcap_\gamma \left(\prod_\beta (I_\gamma E_\beta)\right).$$

Again invoking condition (2) in Lemma 5.2, we may rewrite this intersection as $\cap (I_\gamma P)$, completing the proof. ■

Remark 5.4. Theorem 5.3 (and its proof) may be generalized as follows. Let R be a ring and let $\{E_\beta\}$ be an inverse system of R -modules such that $I \varprojlim E_\beta \cong \varprojlim IE_\beta$ for each finitely generated ideal I of R . Set $E = \varprojlim E_\beta$. Then if R is coherent (resp., Noetherian) and E_β satisfies $(*)$ (resp., $(**)$) for each β , then E satisfies $(*)$ (resp., $(**)$).

This offers an alternate proof of the "if" part of Theorem 4.5. For let (R, M) be quasi-complete. Then R/M^k satisfies $(**)$ over R for each positive integer k . Moreover, for each finitely generated ideal I of R , one has

$$\begin{aligned} I \varprojlim R/M^k &= I\hat{R} \cong I \otimes_R \hat{R} \cong \hat{I} = \varprojlim I/M^k \\ &\cong \varprojlim I/M^k \cap I \cong \varprojlim (I + M^k)/M^k = \varprojlim I(R/M^k). \end{aligned}$$

Thus $\hat{R} = \varprojlim R/M^k$ satisfies $(**)$, as asserted. ■

We next begin the promised reworking of parts of [4].

PROPOSITION 5.5. *Let R be a ring and γ a cardinal number such that each ideal of R has a generating set with cardinality at most γ . Suppose that there exists an R -flat product $P = \prod_\beta R$ of copies of R , with index set having cardinality $\beta \geq \gamma$. Then:*

(a) $I \cap J$ is finitely generated for each pair I, J of finitely generated ideals of R .

(b) $(0:a)$ is finitely generated for each $a \in R$.

Proof. (a) Since P is R -flat, it satisfies $(*)$ by Remark 4.2, whence $(I \cap J)P = IP \cap JP$. Since I and J are each finitely generated, condition (2) of Lemma 5.2 gives $IP = \prod I$ and $JP = \prod J$, all products being indexed by β elements. It follows that $IP \cap JP = \prod (I \cap J)$, whence $(I \cap J) \prod R = \prod (I \cap J)$. Since $\beta \geq \gamma$, the implication $(3) \Rightarrow (1)$ in Lemma 5.2 yields the assertion.

(b) $K = (0:a) = \ker(m_a)$, where $m_a: R \rightarrow R$ is multiplication by a . By flatness of P , $K \otimes_R P$ is identified with $\ker(m_a \otimes \text{id}_P)$; that is, after applying the isomorphism $R \otimes_R P \cong P$, with the kernel of multiplication by a on P .

The upshot is that $K \prod R = \prod K$, each product being indexed by β elements. Since $\beta \geq \gamma$, the implication (3) \Rightarrow (1) in Lemma 5.2 may again be invoked, showing that K is finitely generated, thus completing the proof. ■

COROLLARY 5.6. *An integral domain R is coherent if (and only if) there exist cardinal numbers $\gamma \leq \beta$ such that each ideal of R has a generating set with cardinality at most γ and the product $P = \prod_{\beta} R$ of β copies of R satisfies (*) over R .*

Proof. The “only if” part follows from the criterion in [4, Theorem 2.1(a)] since flatness implies (*). For the “if” part, one need only reread the proof of Proposition 5.5(a) and appeal to the coherence criterion that the intersection of any two finitely generated ideals of R be finitely generated. ■

Our final result gives an alternate proof that products preserve flatness over a coherent integral domain.

PROPOSITION 5.7. *Let R be a (coherent) integral domain such that the intersection of any pair of finitely generated ideals of R is finitely generated. Let $\{E_{\beta}\}$ be a collection of R -flat modules E_{β} . Then $P = \prod E_{\beta}$ is R -flat.*

Proof. Being flat, each E_{β} is torsion-free over R , and hence so is P . Thus, by Proposition 4.1(a), it suffices to prove that P satisfies (*); that is, by the first observation in the proof of Theorem 5.3, that $(I \cap J)P = IP \cap JP$ for all finitely generated ideals I, J of R . Since $I \cap J$ is finitely generated by hypothesis and since each E_{β} (being flat) satisfies (*), we may repeat the calculations displayed in the proof of Theorem 5.3, yielding $(I \cap J)P = (\prod IE_{\beta}) \cap (\prod JE_{\beta})$. Finally, since I and J are finitely generated, Lemma 5.2 reduces this intersection to $IP \cap JP$, completing the proof. ■

REFERENCES

1. D. D. ANDERSON, The existence of dual modules, *Proc. Amer. Math. Soc.* **55** (1976), 258–260.
2. D. D. ANDERSON, On the ideal equation $I(B \cap C) = IB \cap IC$, *Canad. Math. Bull.* **26** (1983), 331–332.
3. N. BOURBAKI, “Commutative Algebra,” Addison-Wesley, Reading, 1972.
4. S. U. CHASE, Direct products of modules, *Trans. Amer. Math. Soc.* **97** (1960), 457–473.
5. D. E. DOBBS, On flat finitely generated ideals, *Bull. Austral. Math. Soc.* **21** (1980), 131–135.
6. D. E. DOBBS, On the criteria of D. D. Anderson for invertible and flat ideals, *Canad. Math. Bull.* **29** (1986), 25–32.

7. R. GILMER, "Finite Element Factorization in Group Rings," Lecture Notes in Pure and Applied Mathematics, Vol. 7, Dekker, New York, 1974.
8. E. W. JOHNSON, A note on quasi-complete local rings, *Colloq. Math.* **21** (1970), 197–198.
9. I. KAPLANSKY, "Infinite Abelian Groups," rev. ed., Univ. of Michigan Press, Ann Arbor, 1969.
10. D. G. NORTHCOTT, "Ideal Theory," Cambridge Tracts in Mathematics and Mathematical Physics, Vol. 42, Cambridge Univ. Press, Cambridge, 1953.
11. F. RICHMAN, Generalized quotient rings, *Proc. Amer. Math. Soc.* **16** (1965), 794–799.
12. H. UDA, LCM-stableness in ring extensions, *Hiroshima Math. J.* **13** (1983), 357–377.